

Supplementary Table 1. Associations between diabetes* and type/frequencies of ACEs stratified by age.

	18-49 years		50 years and older		18-49 years		50 years and older	
	Model 1		Model 2		Model 1		Model 2	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Emotional neglect	1.5	0.9-2.8	1.3	0.9-1.9	1.4	0.8-2.7	1.3	0.9-1.8
Physical neglect	3.4	1.3-9.2	0.8	0.4-1.5	2.8	1.0-8.0	0.9	0.5-1.7
Living with household members who were substance abusers	1.1	0.3-3.4	1.0	0.4-2.2	1.3	0.4-3.9	1.0	0.4-2.4
Living with household members who were mentally ill or suicidal	1.7	0.6-4.8	0.3	0.1-0.6	1.9	0.7-5.4	0.3	0.1-0.6
Battered mother/female guardian	1.5	0.6-3.7	0.9	0.4-2.1	1.4	0.6-3.5	0.8	0.4-1.9
Living with household members who were imprisoned	2.0	0.9-4.9	1.1	0.4-3.0	1.8	0.7-4.5	1.0	0.4-2.6
Parental separation, divorce or death of a parent	2.5	1.3-4.6	1.2	0.8-1.8	2.4	1.3-4.4	1.2	0.8-1.8
Emotional abuse	1.0	0.4-2.6	0.2	0.1-0.5	1.0	0.4-2.7	0.2	0.1-0.4
Physical abuse	2.3	0.9-5.8	0.8	0.3-2.2	2.1	0.8-5.4	0.7	0.2-2.2
Sexual abuse	2.6	0.9-7.5	2.1	0.7-6.9	2.8	0.9-8.4	1.8	0.5-6.5
Bullying	3.2	0.9-11.0	0.3	0.0-1.3	3.7	1.0-13.0	0.3	0.1-1.4
Any ACE	1.5	0.8-2.8	1.3	0.8-1.9	1.4	0.8-2.8	1.3	0.9-2.0
Number of ACEs (Reference=0)								
1	1.1	0.5-2.4	1.2	0.8-2.0	1.1	0.5-2.5	1.3	0.8-2.1
2	1.3	0.5-3.3	1.4	0.8-2.4	1.2	0.5-3.2	1.4	0.8-2.5
3 and more	2.5	1.1-5.5	1.1	0.6-2.1	2.4	1.1-5.5	1.0	0.5-2.0

*Cases with age of onset before 18 years were excluded from the analysis

Model 1: Crude odds ratio (OR);

Model 2: Adjusted OR after controlling for health risk behaviors (i.e., BMI, drinking and smoking)

Significant associations are highlighted in bold